

PHOENIX
TATTOO REMOVAL
← & →
SKIN REVITALIZATION
VASCULAR LESION

Precautions to take before your treatment:

- ☒ No sun exposure, tanning beds and sunless tanning creams, for 4 weeks prior to treatment. Sun exposure decreases the effectiveness of the laser treatment and can increase the chance of post treatment complications.
 - No Antibiotics or/need to be off any antibiotics for at least 3 days.
 - Must be off Accutane at least 6 months
 - No Botox or Fillers of any kind for at least **14 days** prior to treatment.
 - No topical Retin-A or topical acne creams/ointments **3 days prior** to treatment.
 - Remove all makeup, creams, or oils prior to treatment.

VASCULAR LESION POST-TREATMENT GUIDELINES

- A mild sunburn-like sensation is expected. This usually lasts two to twenty-four (2-24) hours but can persist up to seventy-two (72) hours. Mild swelling and/or redness may accompany this, but it usually resolves in two to three (2-3) days. Apply wrapped ice or gel packs to the treatment area for ten to fifteen (10-15) minutes every hour for the next four hours, as needed. Never apply ice directly to skin. An oral, over-the-counter anti-inflammatory (ibuprofen such as Advil®) or an analgesic (acetaminophen such as Tylenol®) may be taken to reduce discomfort. Use medicine according to manufacturer's recommendations.
- The vessels may undergo immediate graying or blanching, or they may exhibit a slight purple or red color change. The vessels will fully or partially fade in about ten to fourteen (10-14) days.
- Until redness has resolved, it is recommended to avoid the following:
 - Applying cosmetics to treated areas.
 - Swimming, especially in pools with chemicals.
 - Hot tubs and Jacuzzis.
 - Activities that cause excessive perspiration.
 - Sun exposure to treated areas. Apply an SPF 45 or greater sunscreen to prevent skin color changes.
 - Aggressive scrubbing and use of exfoliants on the treated area.
- Bathe or shower as usual. Treated areas may be temperature-sensitive.
- Schedule a follow-up appointment for evaluation. Repeat treatments may be performed every seven to fourteen (7-14) days if skin has fully recovered. One to four (1-4) treatments may be necessary seven to fourteen (7-14) days apart.
- Advise persons being treated not to pick, remove, or pull at any darkened lesions as scarring may occur.

VASCULAR LESION TREATMENT POSSIBLE SIDE EFFECTS

- Prolonged itching, redness, and blistering.
- Bruising, peeling, rash, lightening or darkening of skin color, ingrown hairs, purpura, crusting, swelling, infection, and removal or lightening of freckles.
- Removal or lightening freckles.
- Scarring, Risk of incidental hair reduction or removal in the treated areas

Please call or text Judi for any concerns you may have 480-313-4443

****Patient Copy****

PHOENIX TATTOO REMOVAL ← & → SKIN REVITALIZATION

Precautions to take before your treatment:

- ☐ No sun exposure, tanning beds and sunless tanning creams, for 4 weeks prior to treatment. Sun exposure decreases the effectiveness of the laser treatment and can increase the chance of post treatment complications.
 - No Antibiotics or/need to be off any antibiotics for at least 3 days.
 - Must be off Accutane at least 6 months
 - No Botox or Fillers of any kind for at least **14 days** prior to treatment.
 - No topical Retin-A or topical acne creams/ointments **3 days prior** to treatment.
 - Remove all makeup, creams, or oils prior to treatment.

VASCULAR LESION POST-TREATMENT GUIDELINES

- A mild sunburn-like sensation is expected. This usually lasts two to twenty-four (2-24) hours but can persist up to seventy-two (72) hours. Mild swelling and/or redness may accompany this, but it usually resolves in two to three (2-3) days. Apply wrapped ice or gel packs to the treatment area for ten to fifteen (10-15) minutes every hour for the next four hours, as needed. Never apply ice directly to skin. An oral, over-the-counter anti-inflammatory (ibuprofen such as Advil®) or an analgesic (acetaminophen such as Tylenol®) may be taken to reduce discomfort. Use medicine according to manufacturer's recommendations.
- The vessels may undergo immediate graying or blanching, or they may exhibit a slight purple or red color change. The vessels will fully or partially fade in about ten to fourteen (10-14) days.
- Until redness has resolved, it is recommended to avoid the following:
 - Applying cosmetics to treated areas.
 - Swimming, especially in pools with chemicals.
 - Hot tubs and Jacuzzis.
 - Activities that cause excessive perspiration.
 - Sun exposure to treated areas. Apply an SPF 45 or greater sunscreen to prevent skin color changes.
 - Aggressive scrubbing and use of exfoliants on the treated area.

- Bathe or shower as usual. Treated areas may be temperature-sensitive.
- Schedule a follow-up appointment for evaluation. Repeat treatments may be performed every seven to fourteen (7-14) days if skin has fully recovered. One to four (1-4) treatments may be necessary seven to fourteen (7-14) days apart.
- Advise persons being treated not to pick, remove, or pull at any darkened lesions as scarring may occur.

VASCULAR LESION TREATMENT POSSIBLE SIDE EFFECTS

- Prolonged itching, redness, and blistering.
- Bruising, peeling, rash, lightening or darkening of skin color, ingrown hairs, purpura, crusting, swelling, infection, and removal or lightening of freckles.
- Removal or lightening freckles.
- Scarring.
- Risk of incidental hair reduction or removal in the treated areas.

Patient Signature: _____ Date: _____

(Or legal guardian)